Adult CoCM Substance or Alcohol Use Disorder Workflow

Updated: 4/5/2023

screening and assessment

under 18 → See Adolescent Workflow

age 18+

Practice determines how to manage abstinence, risk, and problem use. (see SBIRT for resources).
CoCM starts with dependence

Equivalent to performing the PHQ2

brief screen

positive

negative → reinforce healthy behaviors

Equivalent to performing the PHQ9

AUDIT/DAST/ Other Assess.

abstinence or low risk → reinforce healthy behaviors

high risk or **problem use** → brief intervention

dependence

Determine appropriateness treatment

referral to treatment

Patient

Clinical jud

provider determines enrollment to CoCM

no → Not necessary, requires specialist treatment, emergent, etc. → Practice monitors

yes

Team determines timing for BHCM Assessment. Optional before or after provider visit.

patient agrees

no → Practice monitors

yes → BHCM comprehensive assessment (med, social, beh) status, hx, response

treatment

MAT* | SCR

care coordination and treatment plan

monitor plan*

* see OBAT for OUD protocols

G2086, G2087, G2088 pending time spent on OBAT

Adolescent Substance or Alcohol Use Disorder Workflow

Updated: 4/5/2023

screening and assessment



treatment



* see OBAT for AUD and OUD protocols for adolesents